

Effect of different choices of the Boltzmannized flux operator on thermal exchange and recombination reactions

Koichi Saito*

Tohoku Pharmaceutical University, Sendai 981-8558, Japan

(Dated: February 2, 2008)

The rate constants for recombination and exchange reactions are calculated using the flux correlation approach with a general form of the Boltzmannized flux operator, which can simultaneously describe the Kubo and traditional *half-split* forms. First, we consider an exactly solvable model, i.e., the free particle case, in terms of a new scaling function. Next, as a non-trivial case, we study the recombination and exchange reactions at very high pressure. Since the rate constant is calculated by Laplace transform of the flux correlation function, the result depends on how the Boltzmannized flux operator is chosen. We find that a choice of the flux operator affects the rate constant considerably. For the recombination reaction, the ratio of the rate constant in the *half-split* form to that in the Kubo form approaches zero in the high pressure limit.

I. INTRODUCTION

To study a chemical reaction at the most detailed level, it is necessary to treat the action of the time evolution operator, $\exp(-i\hat{H}t)$ ($\hat{H}$ is the Hamiltonian of the system),[†] onto a (given) initial wave function or density matrix, which is a function of total energy E and total angular momentum J . Such quantum scattering calculations have actually been studied for simple chemical reactions.¹ A time-dependent scattering formalism based on the S -matrix Kohn variational approach² or a coupled channel method in hyperspherical coordinates³ has usually been used to evaluate the quantum reactive scattering cross sections. The number of open vibration-rotation channels, however, increases very rapidly as thermally accessible collision energy becomes high. Therefore, at high energy the exact quantum state-to-state calculations would not be feasible even for a simple reaction. Furthermore, if it is *only* the rate constant that is desired, such a complete calculation for all state-to-state information is *not* economical.

A totally different approach for treating chemical reactions is that based on a correlation between quantum flux operators in the statistical thermodynamics. In the early 60's, Yamamoto⁴ first formulated an exact expression for the rate constant as an application of the general statistical mechanical theory of irreversible process, which was established by Kubo et al.^{5,6} and Mori.⁷ (See also Ref.8.) Independently, Miller et al.^{9,10,11} have developed a method for the rate constant using a time integral of the flux-flux autocorrelation function, which is also exact in the limit that the dynamics is extended to $t \rightarrow \infty$. The feasibility of this approach depends on how to evaluate the time evolution operator for the system, and the correlation is usually calculated using the Feynman path integral technique.^{12,13} Although this method is powerful and convenient to obtain the rate constant, the direct application to large, complicated systems is still difficult because of the notorious sign problem.^{14,15} One approach to improve this situation is a filtering or smoothing technique¹⁶ such as the stationary-phase Monte Carlo.¹⁷ Recently, semiclassical (SC) approaches, implemented via the initial-value representation (SC-IVR), have received a rebirth of interest. (Van Vleck¹⁸ first discussed a drastic approximation based on the semiclassical picture.) A number of studies^{19,20,21} have been carried out along SC-IVR and have demonstrated the capability of these approaches to describe various quantum effects in chemical reactions.

It has also been shown how a quantum mechanical version of the Lindemann mechanism for collisional recombination, where the process is affected by the bath gas M ,



can be handled by the flux-flux autocorrelation function for the $A - B$ collision.^{22,23,24} Some applications of this theory are listed in Refs.25,26. It is furthermore possible to generalize the formalism to include chemical reactions as well as recombination:



[†] We use the natural unit, i.e., $\hbar = c = 1$.

Equations (3) and (4) simultaneously describe the recombination process ($A + BC \rightarrow ABC$) and the exchange reaction ($A + BC \rightarrow AB + C$).²³ This method has been applied to the interesting (combustion) reactions ($O + OH \rightleftharpoons H + O_2$) and the recombination reactions ($O + OH + M \rightarrow HO_2 + M \leftarrow H + O_2 + M$), which are very important in atmospheric chemistry.^{27,28}

As mentioned above, there are two different ways to calculate the rate constant for chemical reactions, i.e., the Yamamoto approach, in which the Kubo form of the flux operator is used, and the Miller approach, in which the traditional *half-split* form is chosen. The two approaches can provide the same result to the rate constant because it is given in terms of the integral of the flux-flux correlation function with respect to time.^{9,11} However, the shapes of the correlation functions calculated by the two approaches are quite different from each other. Therefore, for the recombination and exchange reactions (like Eqs.(3) and (4)) the rate constants in the two approaches could be different, because they are given by Laplace transforms of the flux-flux correlation functions.²⁹ The purpose of this paper is to demonstrate the effect of different choices of the Boltzmannized flux operator on the rate constants for recombination and exchange reactions.

First, we summarize the correlation function method briefly in Sec.II. The expressions of the correlation functions in the Yamamoto and Miller approaches can be unified using a general form of the Boltzmannized flux operator. The difference between the two approaches is discussed explicitly. The rate constants for recombination and exchange processes are also studied. In Sec.III, we consider an exactly solvable model, i.e., the free particle case, in terms of a new scaling function. In Sec.IV, as a non-trivial case, we study the recombination and exchange reactions at very high pressure. Finally, the summary and conclusion are given in Sec.V.

II. FLUX-FLUX CORRELATION APPROACH TO RATE CONSTANTS

The quantum mechanically exact expression for a thermal rate constant $k(T)$ can be written in terms of the flux correlation function^{9†}

$$k(T) = Q_0(T)^{-1} \lim_{t \rightarrow \infty} C_s(r; t), \quad (5)$$

where $Q_0(T)$ is the reactant partition function per unit volume and $C_s(r; t)$ is the *flux-side* correlation function defined by

$$C_s(r; t) = \text{tr}[e^{-\beta \hat{H}} \hat{F}(r) \hat{\mathcal{P}}]. \quad (6)$$

Here, $\hat{F}(r)$ is the *bare* flux operator given by

$$\hat{F}(r) = i[\hat{H}, h(\hat{s})]_{s=r}, \quad (7)$$

with h the Heaviside step function and s the reaction coordinate – see Fig.1. $h(s)$ takes the value of 0(1) in the reactant (product) side of the dividing point r . In Eq.(6), $\hat{\mathcal{P}}$ is the projection operator defined by

$$\hat{\mathcal{P}} = e^{i\hat{H}t} h(\hat{p}) e^{-i\hat{H}t}, \quad (8)$$

with $\hat{p}$ the momentum operator.

For the rate constant, two different approaches have been proposed so far: one is the Yamamoto approach,⁴ which is based on the linear response theory (or the so-called Kubo formula),^{5,6} and the other is the flux-flux autocorrelation function method developed by Miller et al.^{9,10,11} In the Miller approach, Eq.(6) are modified by following two useful facts: the first one is made by noting that $\hat{\mathcal{P}}$ and $\hat{H}$ commute each other (i.e., $[\hat{\mathcal{P}}, \hat{H}] = 0$) in the limit $t \rightarrow \infty$, and the second is to replace $h(\hat{p})$ by $h(\hat{s})$ in Eq.(8), which is also correct in the limit $t \rightarrow \infty$. By virtue of these modifications, the flux-side correlation function in the Miller approach can be rewritten as

$$C_s^M(r; t) = \text{tr}[\hat{F}^M(\beta, r) e^{i\hat{H}t} h(\hat{r}) e^{-i\hat{H}t}], \quad (9)$$

where the superscript M stands for "Miller". Here, $\hat{F}^M(\beta, r)$ is the *half-split* Boltzmannized flux operator

$$\hat{F}^M(\beta, r) = e^{-\beta \hat{H}/2} \hat{F}(r) e^{-\beta \hat{H}/2}. \quad (10)$$

† For simplicity, we consider a reaction in one-dimension.

Using Eq.(7), the rate constant can also be expressed in terms of the *flux-flux* correlation function $C(r, r; t)$ as

$$Q_0(T)k(T) = \int_0^\infty dt C^M(r, r; t), \quad (11)$$

where

$$C^M(r, r; t) = \text{tr}[\hat{F}^M(\beta, r)e^{i\hat{H}t}\hat{F}(r)e^{-i\hat{H}t}]. \quad (12)$$

On the other hand, the linear response theory gives the rate constant

$$Q_0(T)k(T) = \int_0^\infty dt C^Y(r, r; t), \quad (13)$$

where

$$C^Y(r, r; t) = \text{tr}[\hat{F}^Y(\beta, r)e^{i\hat{H}t}\hat{F}(r)e^{-i\hat{H}t}], \quad (14)$$

with

$$\hat{F}^Y(\beta, r) = \frac{1}{\beta} \int_0^\beta d\lambda e^{-(\beta-\lambda)\hat{H}}\hat{F}(r)e^{-\lambda\hat{H}}. \quad (15)$$

Here, the superscript Y stands for "Yamamoto".

If a general form of the Boltzmannized flux operator is introduced by²¹

$$\hat{F}^\kappa(\beta, r) = \frac{1}{\kappa\beta} \int_{(1-\kappa)\beta/2}^{(1+\kappa)\beta/2} d\lambda e^{-(\beta-\lambda)\hat{H}}\hat{F}(r)e^{-\lambda\hat{H}}, \quad (16)$$

where κ is a parameter ($0 \leq \kappa \leq 1$), the correlation functions in the Miller and Yamamoto approaches can be unified as

$$C^\kappa(r, r; t) = \text{tr}[\hat{F}^\kappa(\beta, r)e^{i\hat{H}t}\hat{F}(r)e^{-i\hat{H}t}]. \quad (17)$$

It is easy to check that in the limit $\kappa \rightarrow 0(1)$ Eq.(17) reproduces the correlation function in the Miller (Yamamoto) approach.

Using Eq.(7), the integral with respect to λ in the general form of the Boltzmannized flux operator can be performed:

$$\hat{F}^\kappa(\beta, r) = \frac{i}{\kappa\beta} \left[e^{-(1-\kappa)\beta\hat{H}/2} h(\hat{r}) e^{-(1+\kappa)\beta\hat{H}/2} - e^{-(1+\kappa)\beta\hat{H}/2} h(\hat{r}) e^{-(1-\kappa)\beta\hat{H}/2} \right]. \quad (18)$$

Combining the *partial* Boltzmann operator, $e^{-(1\pm\kappa)\beta\hat{H}/2}$, with the real-time evolution operator $e^{\pm i\hat{H}t}$, we obtain two complex-time evolution operators: $e^{-i\hat{H}t_+}$ and $e^{-i\hat{H}t_-}$ with $t_\pm = t - i(1 \pm \kappa)\beta/2$. Then, the correlation function is rewritten as

$$C^\kappa(r, r; t) = \frac{2}{\kappa\beta} \Im \text{tr}[\hat{F}(r)e^{i\hat{H}t_+^*} h(\hat{r}) e^{-i\hat{H}t_-}], \quad (19)$$

where $\Im$ stands for taking the imaginary part. Performing the trace operation in Eq.(19), the correlation function reads

$$\begin{aligned} C^\kappa(r, r'; t) &\equiv \frac{2}{\kappa\beta} \Im \text{tr}[\hat{F}(r)e^{i\hat{H}t_+^*} h(\hat{r}') e^{-i\hat{H}t_-}] \\ &= \frac{1}{m\kappa\beta} \Im \int_r^\infty ds \left[i \langle s | e^{-i\hat{H}t_+} | s' \rangle^* \frac{\partial}{\partial s'} \langle s | e^{-i\hat{H}t_-} | s' \rangle \right. \\ &\quad \left. - i \langle s | e^{-i\hat{H}t_-} | s' \rangle \frac{\partial}{\partial s'} \langle s | e^{-i\hat{H}t_+} | s' \rangle^* \right]_{s'=r'}, \end{aligned} \quad (20)$$

where $\langle s | e^{-i\hat{H}t} | s' \rangle$ is the propagator in the coordinate representation. The correlation function $C^\kappa(r, r; t)$ is given in the limit $r' \rightarrow r$.

It is possible to generalize the flux correlation approach to treat recombination and exchange reactions.^{22,23,24,25,26} Let us consider the reaction of $A + BC \rightarrow AB + C$ and ABC in one-dimension (see Eqs.(3) and (4) and Fig.2). The rate constants for the exchange ($A + BC \rightarrow AB + C$) and recombination ($A + BC \rightarrow ABC$) reactions are again given by the average of the flux $\hat{F}(r)$ and the projection operator $\hat{P}$ over the Boltzmann distribution, where $\hat{F}(r)$ describes the flux at the reactant dividing point r (see Fig.2). Similarly, we can define the flux operator $\hat{F}(p)$ at the product dividing point p .

Because the probability of the system experiencing a deactivating ($ABC^* + M \rightarrow ABC + M$) collision with the bath gas M can be evaluated by $1 - e^{-\eta t}$ at time t (η describes the frequency of deactivating collisions and it depends on pressure P and T of the bath gas), the recombination probability is proportional to $1 - e^{-\eta\tau}$, where τ is the time the trajectory (it is on r at $t = 0$) is in the compound region (see Fig.2). On the other hand, for the exchange reaction, the probability is given by $e^{-\eta\tau_p}$, where τ_p is the time the trajectory exists through the point p .

Using these probabilities and the correlation function, the rate constants for the recombination and exchange reactions are finally given by²³

$$Q_0(T)k_{rec}^\kappa(T, P) = \int_0^\infty dt e^{-\eta t} [C^\kappa(r, r; t) - C^\kappa(r, p; t)], \quad (21)$$

$$Q_0(T)k_{exc}^\kappa(T, P) = \int_0^\infty dt e^{-\eta t} C^\kappa(r, p; t), \quad (22)$$

where the relation

$$\int_0^\infty dt C^\kappa(r, r; t) = \int_0^\infty dt C^\kappa(r, p; t) \quad (23)$$

holds. This ensures that the recombination rate vanishes in the limit $\eta \rightarrow 0$.

Because the rate constants for recombination and exchange processes are calculated by the Laplace transforms of the correlation functions, it is clear that the evaluated results depend on κ , that is, the shape of the correlation function affects the rate constants.²⁹ Of course, the integral of the correlation function with respect to time itself does not depend on κ .

III. SOLVABLE MODEL – FREE PARTICLE CASE

In this section and the next section, we calculate the rate constants for recombination and exchange reactions using the Feynman path integral technique.^{12,13} A huge calculation is usually required to obtain the exact matrix elements of propagators for a realistic system. Furthermore, it is necessary to take some approximations and numerical techniques to perform it.^{17,19,20,21,30,31,32} Because the aim of this paper is to show how the rate constant for recombination or exchange reaction depends on κ (i.e., a choice of the Boltzmannized flux operator), it would be more intuitive and useful to first consider a simple system rather than a complicated case. We here study an analytically solvable model, i.e., the free particle case in one-dimension, and discuss a non-trivial case in the next section.

If no potential acts on the system (see Fig.3), the propagator in the coordinate representation is easily calculated by the path integral.¹² The flux-flux correlation function Eq.(20) then gives

$$C_0^\kappa(r, p; t) = \frac{1}{4\pi\kappa\beta^2\sqrt{D_+D_-}} \exp\left[-\frac{m\beta d^2(t^2 + (1 - \kappa^2)\beta^2/4)}{2D_+D_-}\right] \\ \times [(2tA - \kappa\beta B) \sin X + (2tB + \kappa\beta A) \cos X], \quad (24)$$

where $D_\pm = t^2 + (1 \pm \kappa)^2\beta^2/4$, $X = \kappa m t \beta^2 d^2 / 2D_+D_-$ and

$$A = \left[(\sqrt{D_+} + t) (\sqrt{D_-} + t) \right]^{1/2} + \left[(\sqrt{D_+} - t) (\sqrt{D_-} - t) \right]^{1/2}, \quad (25)$$

$$B = \left[(\sqrt{D_+} + t) (\sqrt{D_-} - t) \right]^{1/2} - \left[(\sqrt{D_+} - t) (\sqrt{D_-} + t) \right]^{1/2}, \quad (26)$$

with d the distance between r and p and m the reduced mass. Note that, as it should be, the correlation depends on *only* the distance d and is independent of the positions r and p .

If $\kappa = 1$ (the Yamamoto approach), Eq.(24) gives

$$C_0^Y(r, p; t) = \frac{1}{2\pi\beta^2\sqrt{2t(t^2 + \beta^2)}} \exp\left[-\frac{m\beta d^2}{2(t^2 + \beta^2)}\right] \\ \times \left[(\sqrt{t^2 + \beta^2} + t)^{3/2} \sin X' + (\sqrt{t^2 + \beta^2} - t)^{3/2} \cos X' \right], \quad (27)$$

where $X' = m\beta^2 d^2 / 2t(t^2 + \beta^2)$, and

$$C_0^Y(r, r; t) = \frac{(\sqrt{t^2 + \beta^2} - t)^{3/2}}{2\pi\beta^2\sqrt{2t}(t^2 + \beta^2)^{1/2}}. \quad (28)$$

On the other hand, taking the limit $\kappa \rightarrow 0$ in Eq.(24), we obtain the correlation functions in the Miller approach:

$$C_0^M(r, p; t) = \frac{1}{4\pi(t^2 + \beta^2/4)^{3/2}} \left[\frac{\beta}{2} + \frac{2mt^2 d^2}{t^2 + \beta^2/4} \right] \exp \left[-\frac{m\beta d^2}{2(t^2 + \beta^2/4)} \right], \quad (29)$$

and

$$C_0^M(r, r; t) = \frac{\beta}{8\pi(t^2 + \beta^2/4)^{3/2}}. \quad (30)$$

Now it is very convenient to introduce a new scaling function

$$\begin{aligned} S_{rp}^\kappa(x, c) &\equiv 2\pi\beta^2 C_0^\kappa(r, p; t) \\ &= \frac{1}{2\kappa\sqrt{\bar{D}_+\bar{D}_-}} \exp \left[-c \left(\frac{x^2 + (1 - \kappa^2)/4}{\bar{D}_+\bar{D}_-} \right) \right] \\ &\times [(2x\bar{A} - \kappa\bar{B}) \sin X + (2x\bar{B} + \kappa\bar{A}) \cos X], \end{aligned} \quad (31)$$

where $x(=t/\beta)$ and $c(=md^2/2\beta)$ are *dimensionless* variables. Since the scaling function itself is a function of x and c , it is also dimensionless. Here, $\bar{D}_\pm = x^2 + (1 \pm \kappa)^2/4$, $X = \kappa c x / \bar{D}_+\bar{D}_-$, and $\bar{A}$ and $\bar{B}$ are given by replacing $D_\pm$ with $\bar{D}_\pm$ and $t \rightarrow x$ in Eqs.(25) and (26), respectively. If we set $r = p$ (or $d = 0$), we obtain

$$S_{rr}^\kappa(x) \equiv 2\pi\beta^2 C_0^\kappa(r, r; t) = \frac{1}{2\kappa\sqrt{\bar{D}_+\bar{D}_-}} (2x\bar{B} + \kappa\bar{A}), \quad (32)$$

which is a function of *only* x because $c = 0$.

We show these scaling functions in Figs.4 and 5. Note that the integral of the scaling function with respect to x does not, of course, depend on κ . In Fig.4 the scaling function for $\kappa = 1$ (Yamamoto approach) is divergent like $\sim 1/\sqrt{x}$ at $x = 0$. Taking the limit $x \rightarrow 0$ implies that t approaches 0 but β is finite or T is very low ($\beta \rightarrow \infty$) but t is finite. In the Miller approach ($\kappa = 0$), the scaling function is finite at $x = 0$: $S_{rr}^{\kappa=0}(0) = 2$. For arbitrary κ , we find $S_{rr}^\kappa(0) = 2/\sqrt{1 - \kappa^2}$. By contrast, in the limit $x \rightarrow \infty$ (which corresponds to the case where T is very high ($\beta \rightarrow 0$) but t is finite or $t \rightarrow \infty$ but β is finite), the scaling function does not depend on κ and behaves like $\sim 1/4x^3$.

In Fig.5 we take $c = 1.0$ to illustrate the scaling function S_{rp}^κ clearly. It is remarkable that the interference effect in the correlation is fully taken into account in the Yamamoto approach ($\kappa = 1$). Hence the scaling function oscillates very rapidly near the origin. Contrastingly, in the case of $\kappa \neq 1$ the interference is averaged and the shape is very smooth. The value of the scaling function at $x = 0$ is given by $(2/\sqrt{1 - \kappa^2}) \exp(-4c/(1 - \kappa^2))$ for $\kappa \neq 1$. The *partial-split* form of the Boltzmannized flux operator (i.e., $0 \leq \kappa < 1$) is thus very convenient for actual computations. For large x the scaling function is again independent of κ and behaves like $\sim (1 + 8c)/4x^3$.

Next, we study the rate constants for recombination and exchange reactions, which are given by Eqs.(21) and (22), respectively. Using the scaling functions, we first calculate the (rr) - and (rp) -rate constants, which are, respectively, defined by

$$Q_0(T)k_{rr}^\kappa = \frac{1}{2\pi\beta} \int_0^\infty dx e^{-\alpha x} S_{rr}^\kappa(x), \quad (33)$$

$$Q_0(T)k_{rp}^\kappa = \frac{1}{2\pi\beta} \int_0^\infty dx e^{-\alpha x} S_{rp}^\kappa(x, c), \quad (34)$$

with $\alpha(= \beta\eta)$ a dimensionless variable. Then, the rate constants are given as $k_{rec}^\kappa = k_{rr}^\kappa - k_{rp}^\kappa$ and $k_{exc}^\kappa = k_{rp}^\kappa$.

In order to convert the collision frequency η to more familiar variables, we approximate the collisional deactivation rate constant by an expression given by the hard sphere collision theory. Furthermore, if one uses the ideal gas expansion, the frequency can be expressed by²³

$$\eta = k_{deact}[M] = P \sqrt{\frac{2000}{T}} \times 10^{-11}, \quad (35)$$

with η in fs^{-1} , P in Pa and T in K. Then, we find

$$\alpha = \beta\eta \simeq 3.42 \times \frac{P}{T^{3/2}} \times 10^{-6}. \quad (36)$$

The factor c is also converted as

$$c = \frac{md^2}{2\beta} = 0.0103 \times A_m T d^2, \quad (37)$$

with A_m the reduced mass of the system in atomic mass units and d in Å.

Now we are in a position to show our results for the free particle case. We define the ratio of the rate constant for arbitrary κ to that for $\kappa = 1$ (Yamamoto approach):

$$R_{rr}^\kappa(\alpha) = k_{rr}^\kappa / k_{rr}^{\kappa=1}, \quad (38)$$

$$R_{rp}^\kappa(\alpha, c) = k_{rp}^\kappa / k_{rp}^{\kappa=1}. \quad (39)$$

The ratio R_{rr}^κ is a function of *only* α , through which it depends on T and P , while the ratio R_{rp}^κ depends on α and c . In Figs.6 and 7, we illustrate the two ratios for the free particle case. Here, we choose $0 \leq \alpha \leq 1$, which, for example, covers the region of $P \lesssim 1.5$ GPa at $T \sim 300$ K. In Fig.7, c is fixed to be 20 because the ratio is not sensitive to it in the region of $0 \leq \alpha \leq 1$. In R_{rr}^κ , the ratio decreases linearly and it is about 0.9 (10% reduction) at $\alpha = 1$ in the case of $\kappa = 0$. Note that a large value of α corresponds to the case of high P and low T , and that the ratio at large α is mainly determined by the correlation function (or the scaling function) at $x \simeq 0$ (see Fig.4). By contrast, in R_{rp}^κ the deviation of the ratio from unity is not large, and the ratio for $\kappa = 0$ is enhanced by about 4 – 5% at $\alpha = 1$.

Combining the (rr) - and (rp) -rate constants, we can calculate the ratio of the recombination rate constants, $R_{rec}^\kappa = k_{rec}^\kappa / k_{rec}^{\kappa=1}$. Because the (rp) -rate constant itself is, however, much smaller than the (rr) -rate constant, the ratio R_{rec}^κ is very close to R_{rr}^κ . We find that $R_{rec}^{\kappa=0}$ is again about 0.9 at $\alpha = 1$. On the other hand, the ratio for the exchange process is given by $R_{exc}^\kappa = k_{exc}^\kappa / k_{exc}^{\kappa=1} = R_{rp}^\kappa$.

IV. RATE CONSTANTS AT HIGH PRESSURE LIMIT

As a non-trivial example, we consider the rate constants for recombination and exchange reactions at very high pressure. As we have seen in the previous section, it is very notable that at short time (or small x) the correlation function (or the scaling function) in the Yamamoto approach is quite different from the other case ($\kappa \neq 1$). The rate constant is calculated by the Laplace transform of the correlation function (see Eqs.(21) and (22)), in which the integrand involves the factor $\exp(-\eta t)$ and η is proportional to pressure. Thus, at sufficiently high pressure the rate constant may be determined by the correlation function at short time ($t \lesssim \eta^{-1}$).

For a short time, the paths which must be considered in evaluating a potential V never move very far from the initial position. Thus, to a first approximation, we can expand the potential V around the average of the initial (s) and final (s') positions:

$$V(x(t)) \simeq V(\bar{x}) + [x(t) - \bar{x}]V'(\bar{x}) + \frac{1}{2}[x(t) - \bar{x}]^2 V''(\bar{x}), \quad (40)$$

where $\bar{x} = (s + s')/2$. Here we ignore terms of higher order than the second derivative of V . By virtue of the quadratic form of $x(t)$, we can easily find the propagator^{12,13}

$$\langle s' | e^{-i\hat{H}t} | s \rangle = \left[\frac{m\omega}{2\pi i \sin \tau} \right]^{1/2} e^{-itV(\bar{x})} e^{iS}, \quad (41)$$

where

$$S = \frac{m\omega}{4 \sin \tau} (1 + \cos \tau) (s - s')^2 - \frac{f^2 F(\tau)}{2m\omega^3 \sin \tau}, \quad (42)$$

with $\tau = \omega t$, $f = -V'(\bar{x})$, $\omega^2 = V''(\bar{x})/m$ and $F(\tau) = 2(1 - \cos \tau) - \tau \sin \tau$. Because the potential is expanded up to $\mathcal{O}(V'')$, Eq.(41) is valid up to $\mathcal{O}(f^2)$ or $\mathcal{O}(\omega^2)$.

To take the temperature effect into account, it is again necessary to introduce complex-times. Because the correlation function is given by (see Eqs.(16) and (17))

$$C^\kappa(r, p; t) = \frac{1}{\kappa\beta} \int_{(1-\kappa)\beta/2}^{(1+\kappa)\beta/2} d\lambda \operatorname{tr}[e^{-(\beta-\lambda)\hat{H}} \hat{F}(r) e^{-\lambda\hat{H}} e^{i\hat{H}t} \hat{F}(p) e^{-i\hat{H}t}], \quad (43)$$

$$= \frac{1}{\kappa\beta} \int_{-\kappa\beta/2}^{\kappa\beta/2} d\lambda \operatorname{tr}[\hat{F}(r) e^{i\hat{H}(t+i(\beta/2+\lambda))} \hat{F}(p) e^{-i\hat{H}(t-i(\beta/2-\lambda))}], \quad (44)$$

$$\equiv \frac{1}{\kappa\beta} \int_{-\kappa\beta/2}^{\kappa\beta/2} d\lambda C^\kappa(r, p, \lambda; t), \quad (45)$$

we define new complex-times as $t_\pm(\lambda) = t \pm i(\beta/2 \pm \lambda)$. Because of very high pressure (or *short* complex-time), it is clear that the (rp) -rate constant becomes very small compared with the (rr) -rate constant. Therefore, we here focus on the (rr) -rate constant and do not consider the (rp) -rate constant. Then, using the propagator Eq.(41) we obtain

$$C^\kappa(r, r, \lambda; t) \simeq \frac{1}{4\pi[t_+(\lambda)t_-(\lambda)]^{3/2}} \left[\frac{\beta}{2} - \frac{\beta^3}{24m} V''(r) \right] e^{-\beta V(r)}, \quad (46)$$

up to $\mathcal{O}(V'')$. The correlation $C^\kappa(r, r, \lambda; t)$ is completely determined by V and V'' , while the first derivative V' gives a correction to the correlation for the (rp) -rate constant. Because the quantum correction due to the potential does not depend on λ explicitly (see Eq.(46)), the correlation function $C^\kappa(r, r; t)$ can be described in terms of that for the free particle case. It is then written best in the form

$$C^\kappa(r, r; t) = \exp \left[-\beta V(r) - \frac{\beta^2}{12m} V''(r) \right] \times C_0^\kappa(r, r; t). \quad (47)$$

In a typical chemical reaction, the interatomic distance and forces usually range over one or two angstroms, and hence one can expect that the change of the potential is small while the motion of the order of $1\text{\AA}$ has been achieved. Because, in the ideal gas model, the root-mean-square speed of a molecule is usually estimated as $v \sim 1.6\sqrt{T/A_m} \times 10^2$ (m/s), the particle moves by as much as $\Delta x \sim 3.5(T/P\sqrt{A_m}) \times 10^{-4}$ (m) for the short period of η^{-1} . Thus, if $\Delta x \lesssim 1\text{\AA}$, the expansion of the potential Eq.(40) is justified and we can use the correlation function Eq.(47). If we assume a typical mass of $A_m \sim 20$ and $T \sim 300$ K, we find the condition of $P \gtrsim 0.24$ GPa.

Since the correction due to the potential in Eq.(47) is common to the correlation function for arbitrary κ , the ratio of the (rr) -rate constant for arbitrary κ to that in the Yamamoto approach ($\kappa = 1$) is *again* given by the ratio for the free particle case: $R_{rr}^\kappa(\alpha) = k_{rr}^\kappa/k_{rr}^{\kappa=1}$. (Note that $R_{rec}^\kappa \simeq R_{rr}^\kappa$ because at high pressure the (rp) -rate constant is negligibly small.) In Fig.8, we show the ratio R_{rr}^κ in the region of $1 \leq \alpha \leq 10$ (cf. Fig.6). At large α the ratio is reduced considerably, which implies that at very high pressure the (rr) -rate constant for $\kappa \neq 1$ is much smaller than that in the Yamamoto approach.

At high P , the rate constant for $\kappa \neq 1$ may be estimated by

$$Q_0(T)k_{rr}^{\kappa \neq 1} \simeq \frac{Z}{\beta^2} \times S_{rr}^{\kappa \neq 1}(0) \int_0^\infty dx e^{-\alpha x} = \frac{2Z}{\alpha\beta\sqrt{1-\kappa^2}}, \quad (48)$$

where Z is a (dimensionless) constant which involves the correction due to the potential. By contrast, in the case of $\kappa = 1$ the rate constant may be given by

$$Q_0(T)k_{rr}^{\kappa=1} \simeq \frac{Z}{\beta^2} \times [\sqrt{x}S_{rr}^{\kappa=1}(x)]_{x \rightarrow 0} \int_0^\infty dx \frac{e^{-\alpha x}}{\sqrt{x}} = \frac{Z}{\beta} \sqrt{\frac{\pi}{2\alpha}}. \quad (49)$$

The ratio at high pressure is thus given as

$$R_{rr}^\kappa = \frac{2}{\sqrt{1-\kappa^2}} \sqrt{\frac{2}{\pi\alpha}}, \quad (50)$$

which approaches zero in the limit $P \rightarrow \infty$. At $\alpha = 10$ (which, for example, corresponds to the case of $T \sim 300$ K and $P \sim 15$ GPa), Eq.(50) gives $R_{rr}^\kappa = 0.5(0.58)[0.76]$ for $\kappa = 0(0.5)[0.75]$. The value for $\kappa = 0$ very well agrees with the numerical result (see Fig.8), while the value for $\kappa = 0.75$ is a little larger than the numerical one. The approximation Eq.(48) may not work well for large κ because the peak of the scaling function at $x = 0$ becomes sharp near $\kappa = 1$ (see Fig.4). In the present calculation, we find that the (rr) -rate constant in the Yamamoto approach is considerably different from the result calculated by the *partial-split* form. At $\alpha \sim 10$ the rate constant for $\kappa = 1$ is about twice as large as that for $\kappa = 0$.

V. SUMMARY AND CONCLUSION

The exact quantum mechanical expression for thermal reaction rates can be formulated by the linear response theory,^{5,6,7} which Yamamoto first discussed in the early 60's.⁴ Later, Miller et al.^{9,10,11} have independently proposed a more convenient way, i.e., the flux-flux autocorrelation function method, to perform numerical computations. The Miller approach can provide the exact rate constant in the limit that the dynamics of the system is extended to $t \rightarrow \infty$. Using a general form of the Boltzmannized flux operator, we have unified the two approaches and studied the rate constants for thermal exchange and recombination reactions. Because they are calculated by Laplace transforms of the flux-flux correlation functions, the result depends on how to choose the Boltzmannized flux operator.

In this paper, we have first considered a solvable model, i.e., the free particle case, to demonstrate the dependence of the rate constant on κ intuitively. To study it, we have introduced a new scaling function and investigated its properties in detail. As a non-trivial case, we have discussed the (rr) -rate constant at very high pressure. Because under such conditions the reaction rate is determined by a propagator at short time, the ratio of the rate constant in the *partial-split* form of the Boltzmannized flux operator to that in the Yamamoto approach can be described in terms of the scaling function for the free particle case. We have found that the rate constant for recombination reaction in the Yamamoto approach is larger than that in the case of $\kappa \neq 1$. In particular, the (rr) -rate constant in the Yamamoto approach is about twice as large as that in the Miller approach at $\alpha \sim 10$.

In conclusion, the *partial-split* form of the Boltzmannized flux operator is certainly an economical and powerful tool to perform numerical calculations for thermal rate constants of realistic reactions. However, for the recombination reaction it may underestimate the rate constant compared with the result calculated by the linear response theory. The difference could be seen if the experiments for recombination reactions could be performed under the conditions of very high pressure.

* ksaito@tohoku-pharm.ac.jp

- ¹ J. Z. H. Zang, *Theory and Application of Quantum Molecular Dynamics* (World Scientific, Singapore, 1999).
- ² S. L. Mielke, D. G. Truhlar, and D. W. Schwenke, *J. Phys. Chem.* **98**, 1053 (1994).
- ³ A. Kuppermann, *J. Phys. Chem.* **100**, 2621 (1996).
- ⁴ T. Yamamoto, *J. Chem. Phys.* **33**, 281 (1960).
- ⁵ R. Kubo, *J. Phys. Soc. Japan* **12**, 570 (1957).
- ⁶ R. Kubo, M. Yokota, and S. Nakajima, *J. Phys. Soc. Japan* **12**, 1203 (1957).
- ⁷ H. Mori, *J. Phys. Soc. Japan* **11**, 1029 (1956).
- ⁸ P. G. Wolynes, *Phys. Rev. Lett* **47**, 968 (1981).
- ⁹ W. H. Miller, *J. Chem. Phys.* **61**, 1823 (1974).
- ¹⁰ W. H. Miller, *J. Chem. Phys.* **62**, 1899 (1975).
- ¹¹ W. H. Miller, S. D. Schwartz, and J. W. Tromp, *J. Chem. Phys.* **79**, 4889 (1983).
- ¹² R. P. Feynman and A. R. Hibbs, *Quantum Mechanics and Path Integrals* (McGraw-Hill, New York, 1965).
- ¹³ L. S. Schulman, *Techniques and Applications of Path Integrals* (Wiley, New York, 1981).
- ¹⁴ J. D. Doll, D. L. Freeman, and T. L. Beck, *Adv. Chem. Phys.* **78**, 61 (1994).
- ¹⁵ C. H. Mak, R. Egger, and H. Weber-Gottschick, *Phys. Rev. Lett* **81**, 4533 (1998).
- ¹⁶ V. S. Filinov, *Nucl. Phys. B* **271**, 717 (1986).
- ¹⁷ J. D. Doll and D. L. Freeman, *Adv. Chem. Phys.* **73**, 289 (1989).
- ¹⁸ J. V. V. Vleck, *Proc. Natl. Acad. Sci. U.S.A.* **14**, 178 (1928).
- ¹⁹ T. Yamamoto, H. Wang, and W. H. Miller, *J. Chem. Phys.* **116**, 7335 (2002).
- ²⁰ Y. Zhao and W. H. Miller, *J. Chem. Phys.* **117**, 9605 (2002).
- ²¹ T. Yamamoto and W. H. Miller, *J. Chem. Phys.* **118**, 2135 (2003).
- ²² W. H. Miller, *J. Phys. Chem.* **99**, 12387 (1995).
- ²³ T. C. Germann and W. H. Miller, *J. Phys. Chem.* **101**, 6358 (1997).
- ²⁴ W. H. Miller, *J. Phys. Chem.* **102**, 793 (1998).
- ²⁵ J. Qi and J. M. Bowman, *J. Phys. Chem.* **100**, 15165 (1996).
- ²⁶ V. A. Mandelshtam, H. S. Taylor, and W. H. Miller, *J. Chem. Phys.* **105**, 496 (1996).
- ²⁷ J. A. Miller, R. J. Kee, and C. K. Westbrook, *Annu. Rev. Phys. Chem.* **41**, 345 (1990).
- ²⁸ P. O. Wennberg, R. C. Cohen, R. M. Stimpfle, J. P. Koplow, J. G. Anderson, R. J. Salawitch, D. W. Fahey, E. L. Woodbridge, E. R. Keim, R. S. Gao, et al., *Science* **266**, 398 (1994).
- ²⁹ K. Saito, *Bull. Chem. Soc. Jpn.* **76**, 271 (2003).
- ³⁰ C. Leforestier and W. H. Miller, *J. Chem. Phys.* **100**, 733 (1994).
- ³¹ W. H. Thompson and W. H. Miller, *J. Chem. Phys.* **106**, 142 (1997).
- ³² S. X. Sun and W. H. Miller, *J. Chem. Phys.* **117**, 5522 (2002).




FIG. 1: Sketch of a potential surface in one-dimensional reaction versus the reaction coordinate s . The dividing point is denoted by r .




FIG. 2: One-dimensional schematic diagram of the potential for $A + BC \rightarrow AB + C$ reaction. The compound region (ABC) is bounded by the dividing points on reactant (r) and product (p) sides.




FIG. 3: Same as Fig.2, but for the free particle case.




FIG. 4: Scaling function $S_{rr}^{\kappa}(x)$ for the free particle case. The dotted (dot-dashed) [solid] curve is for $\kappa = 0(0.9)[1]$.




FIG. 5: Scaling function $S_{rp}^{\kappa}(x, c)$ for the free particle case. The dotted (dot-dashed) [solid] curve is for $\kappa = 0(0.9)[1]$. We take $c = 1.0$.




FIG. 6: Ratio of the (rr) -rate constants in the free particle case. The solid (dot-dashed) [dotted] curve is for $\kappa = 0(0.5)[0.75]$.




FIG. 7: Same as Fig.6, but for the (rp) -rate constant. The solid (dot-dashed) [dotted] curve is for $\kappa = 0(0.5)[0.75]$. We take $c = 20.0$.




FIG. 8: Ratio of the (rr) -rate constants at very high pressure. The solid (dot-dashed) [dotted] curve is for $\kappa = 0(0.5)[0.75]$.